



ZIKA PREGNANCY GUIDANCE

1. Assess travel history 6 months before pregnancy and during pregnancy for Zika-endemic areas (see CDC website if questionable)
2. Establish dates of travel to endemic country/Symptoms of disease: rash, fever, joint pain, and conjunctivitis
3. If <2 weeks after travel or symptom onset: Zika virus rRT-PCR on serum and urine

For rRT-PCR

- Collect 3 SST (serum separator tubes) of serum and small sterile container of urine and send to ADH lab.
- Label all specimens with patient name and date of birth, collection time and date.
- Complete CDC Form 50.34 and send with specimens.
- Ship cold or on ice - can be refrigerated overnight and carried to local ADH health unit to send to ADH lab by local health unit courier.
- Call local health unit to notify you are sending specimen by courier to ADH lab.

If positive: recent Zika virus infection.

If ≥ 2 weeks after travel or symptoms onset: Zika virus IgM and dengue virus IgM on serum.

See reverse side of this card



ZIKA PREGNANCY GUIDANCE

For IgM

- Collect 3 SST (serum separator tubes) of serum and send to ADH lab.
- Label all specimens with patient name and date of birth, collection time and date.
- Complete CDC Form 50.34 and send with specimens.
- Ship cold or on ice - can be refrigerated overnight and carried to local ADH health unit to send to ADH lab by local health unit courier.
- Call local health unit to notify you are sending specimen by courier to ADH lab.

**Zika virus IgM positive or equivocal and any result on dengue virus IgM:
Presumptive positive**

Call ADH for follow-up CDC confirmation

Prepare to assess infant at birth

Call ADH for guidance

- Daytime: 501-537-8969
- After Hours: 1-800-554-5738

Use standard precautions

***REFER TO CDC UPDATED INTERIM PREGNANCY GUIDANCE**